


ONLINE
AND PRINT

THE CYTOKINE

FactsBook

Katherine A. Fitzgerald
Luke A. J. O'Neill
Andy J. H. Gearing
Robin E. Callard



SECOND
EDITION

THE CYTOKINE *FactsBook*

Second Edition

Katherine A. Fitzgerald

*Department of Biochemistry & Biotechnology Institute,
Trinity College, Dublin, Ireland*

Luke A.J. O'Neill

*Department of Biochemistry & Biotechnology Institute,
Trinity College, Dublin, Ireland*

Andy J.H. Gearing

Biocomm International, Melbourne, Australia

Robin E. Callard

*Institute of Child Health,
University College London, London, UK*



ACADEMIC PRESS

A Harcourt Science and Technology Company

San diego San Francisco New York Boston
London Sydney Tokyo

This book is printed on acid-free paper.

Copyright © 2001 by ACADEMIC PRESS

All Rights Reserved.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher.

Academic Press
A Harcourt Science and Technology Company
Harcourt Place, 32 Jamestown Road, London NW1 7BY, UK
<http://www.academicpress.com>

Academic Press
A Harcourt Science and Technology Company
525 B Street, Suite 1900, San Diego, California 92101-4495, USA
<http://www.academicpress.com>

ISBN 0-12-155142-3

Library of Congress Catalog Number: 2001092474

A catalogue record for this book is available from the British Library

INSTRUCTIONS ON GAINING ACCESS TO THE ON-LINE VERSION OF
THE CYTOKINE FACTSBOOK, SECOND EDITION

Access for a limited period to an on-line version of The Cytokine FactsBook, Second Edition is included in the purchase price of the print edition.

The on-line version has been uniquely and persistently identified by the Digital Object Identifier (DOI)

doi: 10.1006/bkcf.2001

By following the link

<http://dx.doi.org/10.1006/bkcf.2001>

from any Web Browser, buyers of the Cytokine FactsBook, Second edition will find instructions on how to register for access.

If you have any problems with accessing the on-line version, e-mail

tscap@hbtechsupport.com

Typeset by Bibliocraft, Dundee, Scotland
Printed and bound in Great Britain by Bookcraft, Bath, Somerset.
01 02 03 04 05 06 07 BC 9 8 7 6 5 4 3 2 1

Preface
Abbreviations

Sect

Chapter 1
Introduction

Chapter 2
Organisms

Chapter 3
The Cytokine

Chapter 4
Cytokine

Sect

Interleukin

IL-1

IL-2

IL-3

IL-4

IL-5

IL-6

IL-7

IL-8

IL-9

Other

Activin

Amph

Angio

Apo21

APRI

BAFF

4-1BB

BCA-

BDNI

Betac

BMPs

CD27

CD30

CD40

6Ckin

CNTF

CT-1

Contents

Preface	vii
Abbreviations	viii

Section I THE INTRODUCTORY CHAPTERS

<i>Chapter 1</i>	
Introduction	1
<i>Chapter 2</i>	
Organization of the Data	8
<i>Chapter 3</i>	
The Cytokine Network	12
<i>Chapter 4</i>	
Cytokine Receptor Superfamilies	20

Section II THE CYTOKINES AND THEIR RECEPTORS

Interleukins

IL-1	35	IL-10	90
IL-2	44	IL-11	95
IL-3	51	IL-12	99
IL-4	58	IL-13	105
IL-5	64	IL-14	111
IL-6	69	IL-15	113
IL-7	75	IL-16	116
IL-8	80	IL-17	119
IL-9	85	IL-18	123

Other Cytokines and Chemokines (in alphabetical order)

Activin	129	CTACK	200
Amphiregulin	136	EGF	203
Angiostatin	139	ELC	209
Apo2L	142	ENA-78	211
APRIL	148	Eotaxin 1	213
BAFF	150	Eotaxin 2	217
4-1BBL	153	Eotaxin 3	219
BCA-1	157	Epo	221
BDNF	160	Fas ligand	225
Betacellulin	166	FGF	230
BMPs	168	Flt3 ligand	244
CD27L	174	Fractalkine	249
CD30L	178	GCP-2	253
CD40L	183	G-CSF	255
6Ckine	188	GDNF	260
CNTF	192	GITRL	267
CT-1	197	GM-CSF	270

Other names

Lymphocyte-activating factor (LAF), endogenous pyrogen (EP), leukocyte endogenous mediator (LEM), mononuclear cell factor (MCF), catabolin.

THE MOLECULES

Interleukin 1 (IL-1) has a very wide range of biological activities on many different target cell types including B cells, T cells and monocytes¹⁻³. *In vivo*, it induces hypotension, fever, weight loss, neutrophilia and acute phase response. IL-1 α and IL-1 β are distinct molecular forms of IL-1 derived from two different genes. IL-1 α is mostly cell associated and IL-1 β is mostly secreted. Although the amino acid sequence homology between the α and β forms is only about 20%, these molecules bind to the same receptor and have very similar if not identical biological properties. An IL-1 receptor antagonist (IL-1Ra) has been described which is structurally related to IL-1 β and binds to the IL-1 receptor⁴. Intracellular forms of human IL-1Ra have also been identified that are splice variants of IL-1Ra⁵. The antagonist is made by the same cells that secrete IL-1 and may be an important physiological regulator. A cysteine protease (converting enzyme) which releases mature IL-1 β has also been cloned and is termed caspase-1, since it is the founder member of the caspase family of cysteine proteases^{6,7}. A cowpox virus-derived inhibitor (CRMA) of the IL-1-converting enzyme has been shown to inhibit the host inflammatory response⁸.

Crossreactivity

There is 62% amino acid sequence homology between human and mouse IL-1 α and 68% for IL-1 β . Both forms crossreact between humans and mice. There is 77% sequence homology between human and mouse IL-1Ra.

Sources

A wide variety of cells secrete IL-1, including monocytes, tissue macrophages, Langerhans cells, dendritic cells, T lymphocytes, B lymphocytes, natural killer (NK) cells, large granular lymphocytes (LGL), vascular endothelium and smooth muscle, fibroblasts, thymic epithelia, astrocytes, microglia, glioma cells, keratinocytes and chondrocytes.

Bioassays

Activation of murine thymocytes or murine T cell lines. PGE₂ induction in fibroblasts using an IL-1-neutralizing antibody as control. *In vivo* (rabbit) pyrogen assay.

Physicochemical properties of IL-1 α and IL-1 β

Property	IL-1 α		IL-1 β	
	Human	Mouse	Human	Mouse
pI	5	5	7	7
Amino acids – precursor	271	270	269	269
– mature ^a	159	156	153	159
M_r (K) – predicted	18.0	18.0	17.4	17.4
– expressed	17.5	17.4	17.3	17.5
Potential N-linked glycosylation sites ^b	2	3	1	2
Disulfide bonds	0	0	0	0

^a After proteolytic removal of propeptide.

^b IL-1 is not normally glycosylated.

3-D structure

The structure of IL-1 α has been determined at a resolution of 2.7 Å by X-ray crystallography and IL-1 β at lower resolution by NMR spectroscopy. Both forms of IL-1 are stable tetrahedral globular proteins formed by an antiparallel six-stranded barrel closed at one end by a six-stranded β -sheet to form a bowl-like structure.

Gene structure¹¹⁻¹⁴


Scale

Exons 50 aa

Translated

Untranslated

Introns 1Kb



es, Callard,

Other names

Mast cell growth factor (MCGF), multi-colony stimulating factor (multi-CSF), eosinophil-CSF (E-CSF), haematopoietic cell growth factor (HCGF), burst-promoting activity (BPA), P-cell stimulating factor activity, thy-1 inducing factor, WEHI-3 growth factor.

THE MOLECULE

Interleukin 3 (IL-3) is a haematopoietic growth factor which stimulates colony formation of erythroid, megakaryocyte, neutrophil, eosinophil, basophil, mast cell and monocytic lineages¹. IL-3 may also stimulate multipotent progenitor cells, but it is more likely to be important in committing progenitor cells to a differentiation pathway rather than self-renewal of primitive stem cells. Many of the activities of IL-3 are enhanced or depend upon costimulation with other cytokines. IL-3 does not stimulate lymphocyte colony formation, but it is a growth factor for B lymphocytes and it activates monocytes, suggesting that it may have an additional immunoregulatory role. IL-3 has been used clinically to expand haematopoietic precursors after bone marrow transplantation, aplastic anaemia and chemotherapy².

Crossreactivity

Amino acid sequence homology between mouse and human IL-3 is only 29% and there is no cross-species reactivity.

Sources

Activated T cells, mast cells, eosinophils.

Bioassays

Proliferation of TF-1 (human erythroleukaemia), MO7e (human megakaryoblastic leukaemia) or AML-193 (acute myeloid leukaemia) cell lines. Stimulation of erythroid, granulocyte and macrophage colony formation in bone marrow colony assay.

Physicochemical properties of IL-3

Property	Human	Mouse
pI	4-8	4-8
Amino acids – precursor	152	166
– mature ^a	133	140
<i>M_r</i> (K) – predicted	15.1	15.7
– expressed	14-30	28
Potential <i>N</i> -linked glycosylation sites	2	4 ^b
Disulfide bonds	1	2

^a After removal of predicted signal peptide.

^b Glycosylation only at positions 16 and 86 (see sequence). Glycosylation is not required for biological activity.

IL-4

Other names

B cell-stimulating factor 1 (BSF-1).

THE MOLECULE

Interleukin 4 (IL-4) is a pleiotropic cytokine derived from T cells and mast cells with multiple biological effects on B cells, T cells and many nonlymphoid cells including monocytes, endothelial cells and fibroblasts. It also induces secretion of IgG1 and IgE by mouse B cells and IgG4 and IgE by human B cells. The IL-4-dependent production of IgE and possibly IgG1 and IgG4 is due to IL-4-induced isotype switching¹⁻³. IL-4 appears to share this property with IL-13.

Crossreactivity

Two regions of human IL-4 (amino acids 1-90 and 129-149) share approximately 50% sequence homology with the corresponding regions of mouse IL-4. In contrast, the region from amino acid positions 91-128 shares very little homology with the corresponding region of mouse IL-4. There is no cross-species reactivity between human and mouse IL-4.

Sources

Mast cells, T cells, some mouse B-cell lymphomas, bone marrow stromal cells.

Bioassays

Human: Proliferation of PHA T-cell blasts in the presence of blocking anti-IL-2 or anti-IL-2R antibody; proliferation of MO7 cell line; increased expression of CD23 or surface IgM on human tonsillar B cells.

Mouse: Proliferation of CTLL in the presence of anti-IL-2 or anti-IL-2R antibody. Increased expression of MHC class II on murine B cells.

Physicochemical properties of IL-4

Property	Human	Mouse
pI	10.5	6.5
Amino acids – precursor	153	140
– mature ^a	129	120
<i>M_r</i> (K) – predicted	15.0	13.6
– expressed	15-19	15-19
Potential <i>N</i> -linked glycosylation sites	2 ^b	3
Disulfide bonds	3	3

^a After removal of signal peptide.

^b Asn38 is glycosylated.

IL-5

Other names

Eosinophil differentiation factor (EDF), eosinophil colony-stimulating factor (E-CSF), B cell growth factor II (BCGFII), B cell differentiation factor for IgM (BCDF μ), IgA enhancing factor, T cell-replacing factor (TRF).

THE MOLECULE

Interleukin 5 (IL-5) is a T cell-derived glycoprotein which stimulates eosinophil colony formation and is an eosinophil differentiation factor in humans and mice. It is also a growth and differentiation factor for mouse but not human B cells¹⁻³.

Crossreactivity

There is 71% homology between mouse and human IL-5 and significant cross-reactivity in functional assays.

Sources

Mast cells, T cells and eosinophils.

Bioassays

Human: Eosinophil differentiation; proliferation of TF1 cell line.

Mouse: Eosinophil differentiation; proliferation of BCL1 or B13 B-cell lines.

Physicochemical properties of IL-5

Property	Human	Mouse
pI (calculated)	7	7.8
Amino acids – precursor	134	133
– mature ^a	115	113
M_r (K) – predicted	13.1	13.1
– expressed ^b	45	40–50
Potential N-linked glycosylation sites	2	3
Disulfide bonds ^c	2	2

^a After removal of predicted signal peptide.

^b Homodimer.

^c Interchain.

3-D structure

IL-5 is an antiparallel disulfide-linked homodimer. The monomer is biologically inactive. The structure of the dimer has been determined at a resolution of 2.4 Å⁴. It has a novel two-domain structure with each domain showing significant structural homology to the cytokine fold in GM-CSF, M-CSF, IL-2, IL-4 and growth hormone. The IL-5 structure is made up of two left-handed bundles of four α -helices with two short β -sheets on opposite sides of the molecule. The C-terminal strand and helix of one chain of the dimer together with three helices and one strand at the N-terminal end of the other chain make up the bundle of four helices and a β -sheet. This dimeric structure of IL-5 is unique. A 3-D image and PDB file are available from SwissProt P05113.

Other names

Interferon- β 2 (IFN β 2), 26-kDa protein, B cell-stimulatory factor 2 (BSF-2), hybridoma/plasmacytoma growth factor (HPGF or IL-HP1), hepatocyte-stimulating factor (HSF), monocyte granulocyte inducer type 2 (MGI-2), cytotoxic T cell-differentiation factor and thrombopoietin.

THE MOLECULE

Interleukin 6 (IL-6) is a multifunctional cytokine secreted by both lymphoid and nonlymphoid cells which regulates B and T cell function, haematopoiesis and acute phase reactions¹⁻³.

Crossreactivity

There is 42% homology between mouse and human IL-6. Human IL-6 is functional on mouse cells but mouse IL-6 has no activity on human cells.

Sources

IL-6 is made by lymphoid cells (T cells, B cells), and many nonlymphoid cells, including macrophages, bone marrow stromal cells, fibroblasts, keratinocytes, mesangium cells, astrocytes and endothelial cells.

Bioassays

Proliferation by IL-6-dependent B9 cell line. Increased Ig secretion by CESS or other EBV-transformed human lymphoblastoid B cell lines.

Physicochemical properties of IL-6

Property	Human	Mouse
pI (calculated)	6.2	6.5
Amino acids – precursor	212	211
– mature ^a	183 ^b	187
M_r (K) – predicted	20.8	21.7
– expressed	26	22-29
Potential <i>N</i> -linked glycosylation sites	2	0 ^c
Disulfide bonds ^d	2	2 ^d

^a After removal of predicted signal peptide.

^b N-terminal amino acids of human IL-6 derived from a T-cell line, an osteosarcoma cell line and a liposarcoma cell line are Pro, Ala and Val respectively, indicating heterogeneity in the signal peptide cleavage site.

3-D structure

IL-6 has a four antiparallel α -helical structure similar to IL-11, LIF, OSM and GM-CSF. A 3-D image and PDB file are available from SwissProt P05231.